PRELIMINARY REPORT ON SOME PHYSIOLOGICAL PARAMETERS IN PREGNANT SHE CAMEL (Camelus dromedarius) IN NORTH AMERICA

Z.B. Ismail¹, T. Qureshi², M. Levy³, W. Khamas and A. Nour

Department of Basic Medical Sciences, School of Veterinary Medicine, Purdue University, 1240 Lynn Hall, West Lafayette, Indiana 47907, USA ¹Department of Veterinary Clinical Sciences, Faculty of Veterinary Medicine, Jordan University of Science and Technology, Irbid, Jordan ²Fort Dodge Animal Health, P.O.Box 5366, Princeton, New Jersey - 08543 ³Department of Veterinary Clinical Sciences, School of Veterinary Medicine, Purdue University, 1240 Lynn Hall, West Lafayette, Indiana 47907, USA

ABSTRACT

A study was carried out to determine the haematological, serum biochemical enzymes and hormonal values in clinically normal 9 pregnant she camels (*Camelus dromedarius*) aged 8 to 16 years. The mean values of various serum enzymes viz. sorbitol dehydrogenase, aspartate aminotransferase, alkaline phophatase and gamma glutamyltransferase in IU/L were 2.30 ± 0.90 , 71.00 ± 3.0 , 57.0 ± 16.5 and 20.0 ± 2.9 , respectively. The serum levels of various hormones viz. progesterone, cortisol, insulin, tri-iodothyronine and thyroxine were 3.5 ± 1.5 ng/ml, 20.0 ± 5.4 ng/ml, 3.8 ± 1.2 µl/ml, 16.0 ± 3.6 ng/ml and 129.0 ± 16.0 ng/ml, respectively.

Key words: Alkaline phophatase, asparate aminotransferase, cortisol, gamma glutamyltransferase, insulin, pregnant she camel, progesterone, sorbitol dehydrogenase, thyroxine, tri-iodothyronine

Although many studies have been conducted to establish normal haematological, serum biochemical and endocrinological parameters in normal camels, these studies are often incomplete and scattered over many years and in different geographical location in the world (Abdalla et al, 1988; Bengoumi et al, 1997; Ghodasian et al, 1978; Haroun, 1994; Kataria and Bhatia, 1991; Mills and Valli, 1988; Mohamed and Hussein, 1999 and Nazifi and Maleki, 1998). To our knowledge, there are no reports of normal physiological parameters in pregnant she-camels in North America. Therefore, normal haematological, serum biochemical and certain hormonal values in clinically normal pregnant camels were determined.

Materials and Methods

Animals : Nine (8-16 years), clinically healthy, pregnant she-camels (*Camelus dromedarius*) raised on a farm in northeastern Indiana were included in the study. The animals were kept on pasture with a shaded area available and fed alfalfa/

SEND REPRINT REQUEST TO A. NOUR

grass hay mixture and a grain supplement twice a day. The camels were on a regular anthelmintic programme every 3 months. Animals were in the second or third trimester of pregnancy based on the actual breeding dates.

Analyses : Blood samples were aseptically collected from the external jugular vein for haematological and serum biochemical and hormonal analyses. Haematological parameters included red blood cell (RBC) count, packed cell volume (PCV), haemoglobin (Hb), mean corpuscular volume (MCV) and total white blood cell (WBC) count which were analysed using the CELL-DYN 3500 haematology analyser (Abbott Diagnostics, Santa Clara, California, 95054, USA). The mean corpuscular haemoglobin concentration (MCHC) was then calculated according to Mills and Valli formula (Bengoumi *et al*, 1997). The differential white blood cell (WBC) count was determined manually using Giemsa stain.

Serum biochemical parameters included glucose, urea nitrogen, creatinine, phosphorus,

calcium, sodium, potassium, chloride, carbon dioxide, magnesium, triglycerides, cholesterol, sorbitol dehydrogenase (SDH), aspartate aminotransferase (AST), alkaline phosphatase (ALP) and gamma glutamyltransferase (GGT) and their concentrations were measured using a Vitros-System 750 XRC (Johnson and Johnson, USA) based on spectrophotometric principles (Pinus *et al*, 1996). Total serum protein was measured using the biuret method and the various protein fractions were determined using electrophoresis. The albumin/globulin ratio was calculated. Plasma concentrations of fibrinogen were measured using heat precipitation method (McPherson, 1996).

Serum levels of progesterone, cortisol, insulin, thyroxine (T_4) and tri-iodothyronine (T_3) were measured using Immulite (DPC Cirrus Inc., USA) based on radioimmunoassay (RIA) principles (Henry *et al*, 1996).

Results and Discussion

Haematology

The mean values of haematological parameters are presented in table 1. The mean values for RBC and Hb were in agreement with those reported by Al-Ali et al (1988) and Alhadrami (1997) for adult dromedaries racing camels and calves and slightly higher but within the range of those reported by Mohamed et al (1999). The difference in mean values could be due to age, sex, nutritional status and variations in physical and geographical conditions. The MCV and MCHC values were in agreement with previous reported by Alhadrami (1997) and Mohamed et al (1999). Investigations have found that haemato-logical parameters in normal camels do not significantly differ between males and females (Abdalla et al, 1988). The values for total WBC count were similar to the values obtained by Alhadrami (1997) and Mohamed et al (1999) and higher than those reported by Higgins and Kock (1984). The values for segmented neutrophils, lymphocytes, monocytes and eosinophils were slightly higher than those given by Higgins and Kock (1984). Haematological values are comparable with those reported in llamas (Weiser et al, 1992) and cattle (Carlson, 1996). The results of this study showed that camelids have higher neutrophil and lymphocyte counts than cattle

 Table 1. Haematological values in normal pregnant she-camel in North America (n = 9).

Parameters	Mean ± SD	Range
Red blood cells (R BC $10^6/\mu l$)	13.23 ± 1.07	12.0 - 13.8
Packed cell volume (PCV%)	36.61 ± 4.38	29.60 - 42.30
Haemoglobin (Hb g/dl)	15.54 ± 1.82	12.70 - 17.70
Mean corpuscular volume (MCV fl)	32.58 ± 2.40	28.60 - 35.20
Mean corpuscular haemoglobin concentration (MCHC g/dl)	42.51 ± 0.96	41.20 - 44.20
Total white blood cells (WBC $10^3/\mu l$)	13.36 ± 4.19	7.69 - 20.90
Segmented neutrophils $(10^3/\mu l)$	9.25 ± 4.86	4.60 - 17.56
Band neutrophils $(10^3/\mu l)$	0.02 ± 0.05	0.0 - 0.16
Lymphocytes $(10^3/\mu l)$	2.60 ± 1.21	1.15 - 4.46
Monocytes $(10^3/\mu l)$	0.37 ± 0.27	0.0 - 0.92
Eosinophils $(10^3/\mu l)$	1.13 ± 0.60 .	0.21 - 2.36

(Carlson, 1996 and Weiser *et al*, 1992). There is no reported value for band neutrophils in camels in the literature. In this study band neutrophils count was comparable to those reported for New World camelids (Fowler and Zinki, 1989 and Fowler, 1998) and other ruminants (Carlson, 1996). Plasma fibrinogen concentrations in this study are comparable to the values in normal adult dromedaries camels and adult llamas reported by Higgins and Kock (1984) and Fowler and Zinki (1989), respectively.

Serum biochemistry

Mean values of serum urea nitrogen (SUN), creatinine, total serum protein, albumin, globulin, albumin/globulin ratio and total bilirubin in normal North American pregnant she-camels are reported in table 2. SUN value in this report is higher than those given by Abdalla et al (1988); Bengoumi et al (1997) and Mohamed and Hussein (1999). Abdalla et al (1988) opined that hydration status affect the concentration of urea in the blood accounting for the great variation in the normal values. Serum creatinine value was found to be comparable to those values obtained by Abdalla et al (1988) and Mohamed and Hussein (1999) but higher than those obtained by Bengoumi et al (1997). Values of total serum protein and albumin were comparable to those values reported by other researchers (Abdalla et al, 1988; Bengoumi et al 1997; Higgins and Kock, 1984 and Mohamed and Hussein, 1999). Serum globulin value corroborated

those reported by Higgins and Kock (1984) in normal adult dromedaries. Serum value for total bilirubin was similar to those reported by Al-Ali and coworkers (1988) and Mohamed and Hussein (1999). The values reported here for SUN, creatinine, total protein, albumin, globulin and total bilirubin were in agreement to those reported in normal adult llamas (Fowler, 1998) and cattle (Carlson, 1996).

Mean serum concentrations of glucose, triglyceride and cholesterol are reported in table 2. Serum concentrations of glucose in this report was comparable to those obtained by Mohamed and Hussein (1999) but was slightly lower than the results obtained by Al-Ali *et al* (1988); Bengoumi *et al* (1997) and Higgins and Kock (1984). Low blood glucose levels in camels in this study might be due to high demands of glucose metabolism for the developing foetuses. Similar to other New World camelids blood glucose levels in dromedaries were significantly higher than that in other ruminants (Carlson, 1996). High blood glucose levels in camelids can be explained by the

 Table 2. Serum biochemical parameters in normal pregnant she camels (*Camelus dromedarius*) in North America (n=9).

Parameters	Mean ± SD	Range
Urea Nitrogen (SUN) (mg/dl)	28.89 ± 5.51	23.00 - 37.00
Creatinine (mg/dl)	1.75 ± 0.23	1.40 - 2.12
Total serum protein (g/dl)	6.80 ± 0.65	5.80 - 7.90
Albumin (g/dl)	3.30 ± 0.50	2.70 - 3.90
Globulin (g/dl)	3.50 ± 0.30	3.12 - 4.12
Albumin/globulin ratio	0.97 ± 0.12	0.90 - 1.20
Fibrinogen (mg/dl)	300.0 ± 120.0	120.0 - 500.0
Total bilirubin (mg/dl)	0.12 ± 2.30	0.08 – 2.5
Glucose (mg/dl)	120.12 ± 4.90	92.00 - 128.00
Triglycerides (mg/dl)	50.00 ± 20.00	27.00 - 124.00
Cholesterol (mg/dl)	47.00 ± 4.50	45.00 - 58.00
Sodium (mEq/L)	154.67 ± 3.46	169.0 - 160.0
Potassium (mEq/L)	4.67 ± 0.63	4.00 - 5.60
Chloride (mEq/L)	136.00 ± 1.76	131.0 - 137.0
Phosphorus (mg/dl)	5.51 ± 1.13	4.20 - 7.20
Calcium (mg/dl)	9.02 ± 0.38	8.30 - 9.60
Magnesium (mg/dl)	3.12 ± 0.26	2.80 - 3.60
Carbon dioxide (mEq/L)	27.40 ± 3.20	23.0 - 34.0
Anion gap (mEq/L)	17.30 ± 3.50	12.0 - 23.0

Values are means of nine readings from different animals' ± standard deviations.

active gluconeogenesis in these species (Mirgani et al, 1987). The mean concentration of triglycerides in present study was higher than those obtained by Bengoumi et al (1997). This could be due to negative energy balance and mobilisation of lipids during late pregnancy in order to meet energy demands for the growing foetus (Dahlborn et al, 1992). Cholesterol value in pregnant she camels was in agreement with the reported values in adult dromedaries (Bengoumi et al, 1997 and Mohamed and Hussein, 1999) and New World camelids (Carlson, 1996) and was closely similar to that reported for other ruminants (Carlson, 1996). It has been reported that cholesterol concentrations are lower in fit and athletic camels due to reduction in body fat related to regular exercise (Mohamed and Hussein, 1999).

Mean serum concentrations of sodium, potassium, chloride, phosphorus, calcium, magnesium, carbon dioxide and anion gap are presented in table 2. Mean serum concentrations of various electrolytes were similar to those obtained by Abdalla *et al* (1988); Bengoumi *et al* (1997) and Mohamed and Hussein (1999) in adult dromedaries camels and in normal adult llamas and cattle (Carlson, 1996 and Fowler, 1998). However, normal serum values for carbon dioxide and anion gap in dromedary camels were reported for the first time in this study (Table 2).

Serum enzyme activities of sorbitol dehydrogenase (SDH), aspartate aminotransferase (AST), alkaline phosphatase (ALP), and gamma glutamyltransferase (GGT) are reported in table 3. Investigators have observed significant differences in serum enzyme activities in camels kept in various climatic conditions and between different age and sex groups (Bengoumi et al, 1997 and Kataria and Bhatia, 1991). SDH is very specific and sensitive indicator of active hepatocellular damage in domestic animals (Carlson, 1996). Information regarding this enzyme is lacking in the literature on camels. The mean concentration of SDH in this study was similar to the reported values in llamas and adult cattle (Carlson, 1996 and Fowler, 1998). Serum activities of AST and ALP were similar to values obtained by Bengoumi et al (1997) and Mohamed and Hussein (1999) and higher than those obtained by Kataria and Bhatia (1991). They found that serum activities of AST and ALP were significantly higher

during extremely hot conditions and in young male camels. It was concluded according to the researchers that younger animals have less stable cellular membranes resulting in greater leakage of enzymes into the circulation.

Endocrinology

Serum concentrations of various hormones of progesterone, cortisol, insulin, tri-iodothyronine (T_3) and thyroxine (T_4) are given in table 3. There are no comparative studies in the literature of hormonal values in normal pregnant she camel for comparison purposes. The result of progesterone in present study corroborated the findings of Agarwal et al (1992) who reported that the circulating levels of progesterone in pregnant camels fluctuated between 4 and 5 ng/ ml throughout the pregnancy. The animals of this study were pregnant $(2^{nd}/3^{rd}$ trimester) and the reports of Agarwal et al (1992) showed the values at the time of calving. The mean serum insulin concentration in present study was lower than that reported by Dahlborn et al (1992). Nutritional status and timing of sampling could be partly responsible for this discrepancy between the two studies. Thyroid hormones play an important role in the normal development and differentiation of the foetuses in pregnant females (Agarwal et al, 1989). Serum concentrations of thyroxine in pregnant camels in this study were similar to the levels obtained by Agarwal et al (1989) in recently calved camels. However, the reported value for tri-iodothyronine was significantly higher than those obtained by Agarwal et al (1989). This difference in values could be due to one time sampling in this study as well as due to seasonal and climatic variations between studies.

Conclusion

Although the sample size of normal pregnant dromedary camels was small but it provided a complete and comprehensive baseline data of normal haematological, serum biochemical and certain hormonal parameters for pregnant camels.

Acknowledgements

We would like to thank Purdue University for financial assistance and laboratory facilities. We also want to thank Drs. Abdel Salam Talafha and Jennifer Brown for their assistance in

Table 3. Serum activitis of some enzymes and hormones in normal pregnant she camels (*Camelus dromedarius*) in North America (n=9).

Parameters	Mean ± SD	Range
Sorbitol dehydrogenase (SDH) (IU/L)	2.30 ± 0.90	1.00 - 4.00
Aspartate aminotransferase (AST) (IU/L)	71.00 ± 3.00	59.00 - 91.00
Alkaline phosphatase (ALP) (IU/L)	57.00 ± 16.50	45.00 - 83.00
Gamma glutamyltransferase (GGT) (IU/L)	20.00 ± 2.90	16.00 - 24.00
Progesterone (ng/ml)	3.5 ± 1.50	0.20 - 4.7
Cortisol (ng/ml)	20.0 ± 5.40	13.0 - 27.0
Insulin (µl/ml)	3.80 ± 1.20	2.50 - 5.40
Tri-iodothyronine (T ₃) (ng/ml)	16.0 ± 3.60	7.50 - 19.60
Thyroxine (ng/ml) (T_4)	129.0 ± 16.0	87.0 - 125.00

Values are means of nine readings from different animals ± standard deviations.

performing clinical examinations and sample collections.

References

- Abdalla OM, Wasfi IA and Gadir FA (1988). The Arabian race camel normal parameters. 1. Haemogram, enzymes and minerals. Comparative Biochemistry and Physiology 90A:237-239.
- Agarwal SP, Khanna ND, Agarwal VK and Dwaraknath PK (1989). Circulating concentrations of thyroid hormones in pregnant camels (*Camelus dromedarius*). Theriogenology 31:1239-1247.
- Agarwal SP, Rai AK and Khanna ND (1992). Hormonal studies in postpartum female camels and their neonates. Proceedings of Ist International Camel Conference, Dubai. pp 163-168.
- Al-Ali AK, Husayni HA and Power DM (1988). A comprehensive biochemical analysis of the blood of the camel (*Camelus dromedarius*). Comparative Biochemistry and Physiology 89B(1):35-37.
- Alhadrami GA (1997). Comparative haematology in the camel calf and adult racing camel (*Camelus dromedarius*). Journal of Camel Practice and Research 4(1):13.
- Banerjee S adn Bhattacharjee RC (1963). Distribution of body water in the camel (*Camelus dromedarius*). Journal of Physiology 204:1245-1247.
- Bengoumi M, Faye B, De La Farge F, Olson WG and Rico AG (1997). Clinical enzymology in the dromedarian camel (*Camelus dromedarius*). Part I. Enzyme activities and distribution of AST, ALT, GGT, AP and LDH in liver, kidney, muscle, myocardium and blood. Journal of Camel Practice and Research 4(1):19-23.
- Carlson PG (1996). Collection of samples and interpretation of laboratory tests: Clinical chemistry tests. In: Large Animal Internal Medicine. Smith BP (ed.). 2nd Edn. Mosby, St. Louis, Missouri. pp 441-470.

- Dahlborn K, Benlamlih S, Wallsten C and Hossaini-Hilali J (1992). Effects of food deprivation on fluid balance and glucose regulation in pregnant and lactating camels. Proceedings of 1st International Camel Conference, Dubai. pp 275-277.
- Fowler AA and Zinki JG (1989). Reference ranges for haematological and serum biochemical values in llamas (*Lama glama*). American Journal of Veterinary Research 50:2049-2053.
- Fowler ME (1998). Hemic and lymphatic systems. In: Medicine and Surgery of South American Camelids. Fowler, ME (Ed.), 2nd Ed. Iowa State University Press, Iowa. pp 364-376.
- Ghodsian I, Nowrouzian I and Schels HF (1978). A study of some haematological parameters in the Iranian camel. Tropical Animal Health and Production 12:129-132.
- Haroun EM (1994). Normal concentrations of some blood constitutents in young Najdi camels (*Camelus dromedarius*). Comparative Biochemistry and Physiology 128:619-622.
- Henry BJ, Alexander RD and Eng CD (1996). Evaluation of endocrine function. In: Clinical Diagnosis and Management by Laboratory Methods. Henry JB (Ed.), 9th Edn. W.B.Saunders Co., Philadelphia, Pennsylvania. pp 322-373.
- Higgins AJ and Kock RA (1984). The Camel in Health and Disease. 1. A guide to the clinical examination, chemical restraint and medication of the camel. British Veterinary Journal 140:485-504.

- Kataria N and Bhatia JS (1991). Activity of some enzymes in the serum dromedary camels. Research in Veterinary Science 51:174-176.
- McPherson AR (1996). Specific proteins. In: Clinical Diagnosis and Management by Laboratory Methods. Henry JB (Ed.), 9th Ed. W.B.Saunders Co., Philadelphia, Pennsylvania. pp 237-252.
- Mills JN and Valli VO (1988). The haematopoietic system. In: Robinson, WF and Huxtable CRC (Eds.). Clinicopathologic principles for Veterinary Medicine. Cambridge University Press, Cambridge. pp 33-84.
- Mirgani T, Uro AO and Sallman HP (1987). Studies on gluconeogenesis in the camel. University of Khartoum, Fac. Vet. Sci. Report Part I. pp 62-70.
- Mohamed HA and Hussein AN (1999). Studies on normal haematological and serum biochemical values of the 'Hijin' racing camels (*Camelus dromedarius*) in Kuwait. Veterinary Research Communication 23:241-248.
- Nazifi S and Maleki K (1998). Biochemical analysis of serum and cerebrospinal fluid in clinically normal adult camels (*Camelus dromedarius*). Research in Veterinary Science 65:83-84.
- Pinus MR, ZImmerman HJ and Henry JB (1996). Clinical enzymology. In: Clinical Diagnosis and Management by Laboratory Methods. Henry JB (Ed.), 9th Edn. W.B. Saunders sCo., Philadelphia, Pennsylvania. pp 268-295.
- Weiser MG, Fettman MJ, Van Houten D, Johnson L and Garry F (1992). Characterisation of erythrocytic indices and serum iron values in healthy llamas. American Journal of Veterinary Research 53(10):1776-1779.